



**The Effect of Acupressure and Foot Hydrotherapy Using Red Ginger
(*Zingiber officinale var rubrum*) on the Blood Pressure of Pregnant
Women with Preeclampsia at Kembaran I Health Center**

Ratna Sulistyana¹, Choiruel Anwar², Walin³

^{1,3}Department of Midwifery, Poltekkes Kemenkes Semarang, Indonesia

²Department of Environmental Health, Poltekkes Kemenkes Semarang, Indonesia

¹Jl. Tirto Agung Pedalangan Banyumanik, Semarang, Jawa Tengah, Indonesia

^{2,3}Jl. Raya Baturraden KM.12, Karangmangu, Banyumas, Jawa Tengah, Indonesia

Corresponding author: Ratna Sulistyana

Email: ratnamedica2@gmail.com

Received: written by editor; Revised: written by editor; Accepted: written by editor

ABSTRACT

The research problem is to determine the effect of Acupressure and Foot Hydrotherapy interventions using red ginger (*Zingiber officinale var rubrum*) on the blood pressure of pregnant women with preeclampsia at Kembaran I Health Center. This study employs a quantitative experimental method with a total of 15 pregnant women with preeclampsia as research subjects. The sampling technique used is non-probability sampling with inclusion criteria of pregnant women with preeclampsia, a gestational age of ≥ 20 weeks, and a Mean Arterial Pressure (MAP) value of >90 mmHg. The measurement instruments consist of a digital sphygmomanometer to measure blood pressure before and after the intervention, and an observation sheet to record pre- and post-intervention blood pressure (MAP) values. Data analysis is conducted using Paired Samples Correlations and paired sample t-tests. The analysis results reveal that the Acupressure and Foot Hydrotherapy therapy using red ginger has a significant effect on lowering blood pressure in pregnant women with preeclampsia. This is evidenced by the T-test value being greater than the t-table value at a 5% alpha level and degrees of freedom (n-1), which is (2.145). This indicates a significant difference between pre- and post-intervention blood pressure after the Acupressure and Foot Hydrotherapy using red ginger. Furthermore, the analysis of Paired Samples Correlations shows a very high correlation between pre- and post-intervention systolic pressure, diastolic pressure, and MAP. This demonstrates a strong correlation between pre- and post-intervention values.

Keyword: *Acupressure, Foot Hydrotherapy, Red Ginger, Blood Pressure, and Preeclampsia*

Introduction

The Maternal Mortality Rate (MMR) in Indonesia in 2023 recorded 4,482 deaths, representing a 25.48% increase compared to 3,572 deaths in 2022. The leading cause of maternal death in 2023 was preeclampsia, with 412 cases, followed by hemorrhage with 360 cases, and 204 other cases related to obstetric complications. Data indicates that preeclampsia is a major contributor to the high maternal mortality rate in Indonesia¹.

However, the Maternal Mortality Rate in Banyumas Regency has shown a significant

downward trend over the past three years. In 2021, there were 44 maternal deaths, which decreased to 24 cases in 2022, and further reduced to 19 cases in 2023. Analysis of the etiological factors contributing to maternal deaths revealed that congenital diseases, brain neoplasms, and chronic liver conditions each contributed to 6 cases of death. Preeclampsia and hemorrhagic events each accounted for 5 cases, while cardiovascular disorders, tuberculosis, and amniotic fluid embolism each contributed 1 case. In 2023, preeclampsia and hemorrhage ranked as the second

highest causes of maternal mortality in Banyumas Regency, comprising 10 cases (52.63%) of MMR. Based on this phenomenon, preeclampsia and hemorrhagic events are the primary causes of maternal death in Banyumas Regency².

Preeclampsia is a pregnancy disorder characterized by elevated blood pressure with systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg, which appears after 20 weeks of gestation and often occurs toward the end of pregnancy. Preeclampsia is classified into mild and severe types based on clinical signs, laboratory results, and the presence of complications. Severe preeclampsia is defined as systolic blood pressure reaching ≥ 160 mmHg or diastolic blood pressure ≥ 110 mmHg, measured on two separate occasions with at least 4 hours apart. Preeclampsia is always followed by new onset hypertension with proteinuria, which indicates the development of hypertension and proteinuria during pregnancy. Preeclampsia can be diagnosed even without proteinuria, but may be accompanied by thrombocytopenia ($<100,000$), elevated SGOT/SGPT (twice the normal levels), right upper abdominal pain, shortness of breath, pulmonary edema, serum creatinine > 1.1 mg/dL, neurological symptoms, visual disturbances, headache, stroke, and fetal growth disturbances⁴.

In the diagnosis of **Severe Preeclampsia (PEB)**, in addition to elevated systolic blood pressure ≥ 160 mmHg and diastolic blood pressure ≥ 110 mmHg, thrombocytopenia (platelet count $< 100,000/\text{microliter}$), liver dysfunction (abnormal or twice the normal liver enzyme concentration), epigastric pain, progressive renal insufficiency (serum creatinine > 1.1 mg/dL or double the normal levels in the absence of kidney disease), and the presence of edema and visual disturbances are also considered⁴.

Various efforts made by the government to reduce **MMR (Maternal Mortality Rate)** aim to ensure quality healthcare services for all pregnant women, including antenatal care, skilled assistance during labor, postpartum support for mothers and babies, providing special care, referring cases with complications, and offering postpartum family planning services at healthcare facilities¹. Efforts to prevent preeclampsia represent a significant advancement in prenatal care, categorized systematically into primary, secondary, and tertiary prevention⁴. Primary prevention is the most effective and can be achieved by mitigating risk factors that may trigger the condition. Secondary prevention aims to prevent the progression of

preeclampsia before clinical manifestations occur, while tertiary prevention focuses on mitigating complications arising as a consequence of the disease⁴.

Primary care requires healthcare professionals capable of identifying risk factors for preeclampsia according to guidelines outlined in the revised 2022 KIA (Kesehatan Ibu dan Anak) book, which includes screening for risk factors early in pregnancy before 20 weeks of gestation. This screening includes assessing parity, maternal age, previous preeclampsia occurrences, BMI evaluation, Mean Arterial Pressure (MAP) assessment, and laboratory testing using urinary dipstick for protein as one of the efforts to prevent preeclampsia. If one or more risk factors for preeclampsia are identified through screening, a low-dose aspirin (75 mg/day) and calcium supplementation (1-1.5 grams) are recommended, and pregnant women with preeclampsia are advised to regularly monitor their blood pressure with a local midwife every 3 days. According to the Family Health Report from Kembaran I Health Center in 2023, there were 98 cases of pregnant women with preeclampsia⁶.

The high incidence of preeclampsia at Kembaran I Health Center prompted healthcare providers to innovate in addressing the issue using a non-pharmacological approach that combines modern and traditional therapeutic elements, which are easy to apply independently and use materials readily available in the community, such as acupressure and hydrotherapy. Several studies on non-pharmacological therapies, such as acupressure, have shown significant differences in average systolic blood pressure before and after acupressure therapy for hypertension in pregnant women during the second trimester⁷. Acupressure is a physiotherapy modality that involves applying massage and stimulation to specific acupoints on the body. Meanwhile, Widiastini states that hydrotherapy, with warm water immersion and foot massage, can reduce edema in the lower extremities of pregnant women⁸. Applying warm foot baths (hydrotherapy) is effective in lowering blood pressure in pregnant women because warm water helps dilate blood vessels and improve blood circulation⁹.

Non-Pharmacological Therapy to address the risk of hypertension in pregnant women with preeclampsia will include **Foot Hydrotherapy** combined with **Red Ginger (Zingiber Officinale var rubrum)**, as red ginger is widely available in Indonesia and provides a spicy sensation along with

the characteristic aroma of ginger from essential oils such as zingiberene, zingiberol, bisabolene, curcumin, gingerol, philandrene, and bitter resin¹⁰. Red ginger contains the highest essential oil content compared to other varieties, which is 2.58-3.90%, while white ginger has 1.5–3.5%, and yellow ginger has the lowest essential oil content, 0.82–1.66%¹¹. The spicy sensation and the warmth from the essential oils in red ginger help dilate blood vessels, improve blood circulation, and ease the heart's workload, which results in lowered blood pressure¹².

Based on this phenomenon, the researcher is interested in conducting a study on the Effect of Acupressure and Foot Hydrotherapy Using Red Ginger (*Zingiber Officinale* var *rubrum*) on blood pressure in pregnant women with preeclampsia at Kembaran I Health Center, Banyumas.

Methods

This study is a quantitative experimental research aimed at analyzing the effect of acupressure and foot hydrotherapy using red ginger (*Zingiber Officinale* var *rubrum*) on blood pressure in pregnant women with preeclampsia at Kembaran I Health Center. The study subjects are pregnant women with preeclampsia during the period of October-November 2024, totaling 15 respondents, with a non-probability sampling technique.

The inclusion criteria for this study are pregnant women with preeclampsia with a gestational age ≥ 20 weeks and a Mean Arterial Pressure (MAP) > 90 mmHg. The measurement tools include a digital sphygmomanometer (OMRON brand) for measuring blood pressure, certified with calibration certificate number 11-31-83309, an observation sheet for recording blood pressure (MAP) before and after the intervention, and a daily schedule sheet to record the implementation of Acupressure and Foot Hydrotherapy with red ginger, which will be performed 6 times over 2 weeks.

Data analysis will include univariate analysis (calculating numerical data, presented as mean, standard deviation, maximum and minimum values with a 95% confidence interval) and bivariate analysis (assessing normality and homogeneity using Levene's test and Shapiro-Wilk test). The difference in mean pre- and post-intervention will be evaluated using paired sample t-test. The study has received ethical approval from the Health Research Ethics Committee of Poltekkes Kemenkes

Semarang, with approval number No. 1231/EA/F.XXIII.38/2024.

Results and Discussion

Univariate Analysis

The results of the descriptive statistical analysis of the blood pressure of pregnant women with preeclampsia at Kembaran I Health Center before and after the intervention are presented in Table 1.

Table 1. Descriptive Statistical Analysis of Blood Pressure

	Min	Max	Mean	Std. Dev
PreSistol_1	112.00	141.00	124.80	8.8898
PostSistol_1	101.00	127.00	111.33	6.3546
PreDiastol_1	78.00	90.00	83.73	3.1275
PostDiastol_1	67.00	87.00	76.73	5.0915
PreMAP_1	91.00	105.00	97.06	4.1248
PostMAP_1	80.60	97.00	88.17	4.7762
PreSistol_2	108.00	137.00	120.27	7.8692
PostSistol_2	102.00	129.00	110.40	7.4718
PreDiastol_2	67.00	95.00	80.33	7.3840
PostDiastol_2	65.00	90.00	73.87	6.7704
PreMAP_2	80.60	105.00	93.51	6.6922
PostMAP_2	78.30	99.00	85.98	6.3821
PreSistol_3	100.00	133.00	119.00	9.0947
PostSistol_3	95.00	121.00	110.80	7.4757
PreDiastol_3	67.00	95.00	79.40	8.1574
PostDiastol_3	66.00	93.00	74.73	6.9843
PreMAP_3	80.60	104.00	92.50	7.3149
PostMAP_3	78.30	102.00	86.63	6.2590
PreSistol_4	107.00	138.00	120.67	8.5161
PostSistol_4	99.00	124.00	109.93	6.8187
PreDiastol_4	66.00	90.00	79.733	7.4495
PostDiastol_4	66.00	81.00	74.73	5.3648
PreMAP_4	80.00	102.00	93.44	6.9317
PostMAP_4	77.00	93.30	86.44	5.1652
PreSistol_5	104.00	142.00	118.00	8.6355
PostSistol_5	97.00	126.00	109.07	6.5188
PreDiastol_5	66.00	96.00	78.93	7.6201
PostDiastol_5	64.00	82.00	74.13	5.3434
PreMAP_5	81.30	104.00	91.89	6.4870
PostMAP_5	77.00	93.30	85.70	4.5551
PreSistol_6	108.00	134.00	116.00	7.1514
PostSistol_6	101.00	114.00	107.27	4.0789
PreDiastol_6	67.00	88.00	77.40	6.1505
PostDiastol_6	65.00	84.00	72.53	6.1860
PreMAP_6	81.30	99.30	90.23	5.3887
PostMAP_6	77.00	92.00	84.09	4.6951

Source: Data processed, 2024

Based on Table 1, it can be seen that the blood pressure before and after the intervention shows a significant decrease. The intervention of acupressure and foot hydrotherapy using red ginger

in 15 pregnant women with preeclampsia for two weeks with six interventions resulted in a decrease in blood pressure. On the first day, the average pre-systolic pressure was 124.80 mmHg (standard deviation 8.8898), while the post-systolic pressure was 111.33 mmHg (standard deviation 6.3546), a decrease of 13.47 mmHg. For pre-diastolic pressure, the average was 83.73 mmHg (standard deviation 3.1275), which decreased to 76.73 mmHg (standard deviation 5.0915), a reduction of 7 mmHg. The mean arterial pressure (MAP) pre-intervention was 97.06 mmHg (standard deviation 4.1248), which decreased to 88.17 mmHg (standard deviation 4.7762), a decrease of 8.89 mmHg.

For the subsequent days, over the course of 2 weeks with 6 interventions, the average pre-systolic values for the minimum, maximum, and mean were higher than the post-systolic values for the minimum, maximum, and mean. Similarly, the average pre-diastolic values for the minimum, maximum, and mean were higher than the post-diastolic values. The average pre-MAP values for the minimum, maximum, and mean were also higher than the post-MAP values.

Bivariate Analysis

Normality Test

The results of the normality test using the Kolmogorov-Smirnov and Shapiro-Wilk tests are shown in Table 2 below.

Tabel 2. Data Normality Test

Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Stat	df	Sig.	Stat	df	Sig.
Pre intervensi	0.127	45	0.068	0.934	45	0.013
Post intervensi	0.137	45	0.033	0.923	45	0.005

Sumber : Data diolah, 2024

Based on Table 2, it can be seen that the pre-intervention test showed a significance value (sig) less than the cut-off value of 5%, indicating that the data distribution was normal. However, the other variables did not follow a normal distribution as their significance values were less than 5%. Therefore, data transformation was performed for re-analysis. The results of the data analysis based on the transformed data, using the Kolmogorov-Smirnov and Monte Carlo tests, are presented in Table 3.

Based on Table 3, the Kolmogorov-Smirnov value for pre-intervention was 0.776 with an Asymp. sig (2-tailed) value of 0.584 (> 0.05). After the intervention, the Kolmogorov-Smirnov value was 0.767 with an Asymp. sig (2-tailed) value of 0.599 (> 0.05). This indicates that the data followed a normal distribution. The results of the Monte Carlo analysis showed a Sig (2-tailed) value of 0.558 for pre-intervention (> 0.05), and for post-intervention, the Sig value was 0.571 (> 0.05). Therefore, it can be concluded that the research data followed a normal distribution.

Tabel 3. Normality Test Based on Transformed Data

One-Sample Kolmogorov-Smirnov Test						
		Sqrtpre		Sqrtpost		
N		90		90		
Normal Parameters ^{a,b}	Mean	10.1782		9.8609		
	Std. Deviation	.99789		.98791		
	Absolute	.082		.081		
Most Extreme Differences	Positive	.082		.081		
	Negative	-.061		-.049		
Kolmogorov-Smirnov Z		.776		.767		
Asymp. Sig. (2-tailed)		.584		.599		
Monte Carlo Sig. (2-tailed)	Sig.	.558 ^c		.571 ^c		
	99% Confidence Interval	.546		.558		
	Lower Bound	.546		.558		
	Upper Bound	.571		.584		

a. Test distribution is Normal.

b. Calculated from data.

c. Based on 10000 sampled tables with starting seed 2000000.

Source: Data processed, 2024

Homogeneity Test

The results of the homogeneity test for pre and post-intervention of Acupressure and Foot Hydrotherapy using red ginger on pregnant women with preeclampsia at Puskesmas Kembaran I are presented in Table 4.

Tabel 4. Homogeneity Test

		Levene Statistic	df ₁	df ₂	Sig.
Pre	Based on mean	2,365	1	88	0.128
	Based on median	1.269	1	88	0.263
	Based on median & with adjusted df	1.269	1	83.4	0,263
	Based on trimmed mean	2.317	1	88	0,132
Post	Based on mean	4.171	1	88	0,044
	Based on median	3.234	1	88	0,076
	Based on median & with adjusted df	3.234	1	83.0	0,076
	Based on trimmed mean	4.048	1	88	0,047

Source: Data processed, 2024

Based on Table 4, it can be seen that for the pre-intervention, the significance values are as follows; Pre based on mean (0.128), Pre based on median (0.263), Pre based on median and with

adjusted df (0.263), Pre based on trimmed mean (0.132). These significance values are > 0.05 , indicating homogeneity. After the intervention, the significance values are; Post based on mean (0.044), Post based on median (0.076), Post based on median and with adjusted df (0.076), Post based on trimmed mean (0.047). Since some significance values are < 0.05 , this indicates a lack of homogeneity. Therefore, data transformation was applied. The results of the pre-intervention analysis with the transformed data are presented in Table 5.

Tabel 5. Homogeneity Test for Pre-Intervention Based on Transformed Data

Levene's Test of Equality of Error Variances ^a			
Dependent Variable: Sqrtpre			
F	df1	df2	Sig.
1.014	1	88	0,317
Tests the null hypothesis that the error variance of the dependent variable is equal across groups.			
a. Design: Intercept + kelompok			
Source: Data processed, 2024			

Based on Table 5, it can be seen that for the pre-intervention, the Levene's test of equality of error variances resulted in an F value of (1.014) with a significance value of 0.317 (> 0.05). This indicates that homogeneity occurred. Meanwhile, the post-intervention Levene's test of equality of error variances is presented in Table 6.

Tabel 6. Homogeneity Test for Post-Intervention Based on Transformed Data

Levene's Test of Equality of Error Variances ^a			
Dependent Variable: Sqrtpost			
F	df1	df2	Sig.
2.108	1	88	0,150
Tests the null hypothesis that the error variance of the dependent variable is equal across groups.			
a. Design: Intercept + kelompok			
Source: Data processed, 2024			

Based on Table 6, the results of the post-intervention homogeneity test using Levene's test of equality of error variances showed an F value of (2.108) with a significance value of 0.150 (> 0.05). This indicates that homogeneity occurred.

Blood Pressure Measurement (MAP)

Based on the results of the Paired Samples Correlations analysis for pregnant respondents with preeclampsia after being intervened with

Acupressure and Foot Hydrotherapy using red ginger at Puskesmas Kembaran I, the results can be seen in Table 7 below.

Tabel 7. Paired Samples Correlations Analysis

		Correlation	Sig.
Pair 1	PreSistol_1 & PostSistol_1	0.827	0.000
Pair 2	PreDiastol_1 & PostDiastol_1	0.323	0.241
Pair 3	PreMAP_1 & PostMAP_1	0.708	0.003
Pair 4	PreSistol_2 & PostSistol_2	0.783	0.001
Pair 5	PreDiastol_2 & PostDiastol_2	0.814	0.000
Pair 6	PreMAP_2 & PostMAP_2	0.866	0.000
Pair 7	PreSistol_3 & PostSistol_3	0.780	0.001
Pair 8	PreDiastol_3 & PostDiastol_3	0.867	0.000
Pair 9	PreMAP_3 & PostMAP_3	0.842	0.000
Pair 10	PreSistol_4 & PostSistole_4	0.714	0.003
Pair 11	PreDiastol_4 & PostDiastol_4	0.590	0.021
Pair 12	PreMAP_4 & PostMAP_4	0.710	0.003
Pair 13	PreSistol_5 & PostSistol_5	0.921	0.000
Pair 14	PreDiastol_5 & PostDiastol_5	0.844	0.000
Pair 15	PreMAP_5 & PostMAP_5	0.889	0.000
Pair 16	PreSistol_6 & PostSistol_6	0,629	0,012
Pair 17	PreDiastol_6 & PostDiastol_6	0.814	0.000
Pair 18	PreMAP_6 & PostMAP_6	0.815	0.000

Source: Data processed, 2024

Based on Table 7, it can be seen that post-intervention Paired Samples Correlations for pre and post systolic, pre and post diastolic, as well as pre and post MAP from the Acupressure and Foot Hydrotherapy intervention using red ginger for pregnant women with preeclampsia showed a relatively high correlation with a significance value of < 0.05 . This indicates a significant correlation between pre and post systolic, pre and post diastolic, as well as pre and post MAP.

This means that after the intervention of Acupressure and Foot Hydrotherapy using red ginger, there was a significant decrease in blood pressure among pregnant women with preeclampsia. However, there was one paired data between pre and post diastolic showing a relatively low correlation of 0.323 with a significance value of 0.241. This indicates a weak relationship between pre and post diastolic.

The results of the Paired Samples T Test analysis for pregnant women with preeclampsia after being intervened with Acupressure and Foot Hydrotherapy using red ginger at Puskesmas Kembaran I are presented in Table 8 below.

Tabel 8 Paired Samples T-Test Analysis

		Paired Differences					t	df
		Mean	Std. Dev	Std. error mean	95% confidence interval of the difference			
					Lower	Upper		
Pair 1	PreSistol_1 - PostSistol_1	13.467	5.097	1.316	10.644	16.289	10.232	14
Pair 2	PreDiastol_1 - PostDiastol_1	7.000	5.043	1.302	4.207	9.792	5.376	14
Pair 3	PreMAP_1 - PostMAP_1	8.887	3.453	0.892	6.974	10.799	9.966	14
Pair 4	PreSistol_2 - PostSistol_2	9.867	5.069	1.309	7.059	12.674	7.539	14
Pair 5	PreDiastol_2 - PostDiastol_2	6.467	4.357	1.125	4.054	8.879	5.749	14
Pair 6	PreMAP_2 - PostMAP_2	7.533	3.395	.87650	5.653	9.413	8.595	14
Pair 7	PreSistol_3 - PostSistol_3	8.200	5.709	1.474	5.038	11.362	5.562	14
Pair 8	PreDiastol_3 - PostDiastol_3	4.667	4.065	1.049	2.415	6.918	4.446	14
Pair 9	PreMAP_3 - PostMAP_3	5.867	3.942	1.018	3.683	8.049	5.764	14
Pair 10	PreSistol_4 - PostSistol_4	10.733	6.006	1.551	7.407	14.059	6.922	14
Pair 11	PreDiastol_4 - PostDiastol_4	5.000	6.095	1.5734	1.625	8.375	3.177	14
Pair 12	PreMAP_4 - PostMAP_4	7.000	4.885	1.262	4.295	9.705	5.550	14
Pair 13	PreSistol_5 - PostSistol_5	8.933	3.654	.943	6.909	10.957	9.468	14
Pair 14	PreDiastol_5 - PostDiastol_5	4.800	4.229	1.092	2.458	7.142	4.396	14
Pair 15	PreMAP_5 - PostMAP_5	6.187	3.205	.827	4.412	7.961	7.477	14
Pair 16	PreSistol_6 - PostSistol_6	8.733	5.574	1.439	5.647	11.819	6.068	14
Pair 17	PreDiastol_6 - PostDiastol_6	4.867	3.758	.970	2.785	6.948	5.015	14
Pair 18	PreMAP_6 - PostMAP_6	6.140	3.135	.809	4.404	7.876	7.586	14

Sumber: data diolah, 2024

Based on Table 8, it can be seen that the T-Test values for pre and post systolic, pre and post diastolic, as well as pre and post MAP during 2 weeks with 6 interventions of Acupressure and Foot Hydrotherapy using red ginger showed that the T-Test values are greater than the t-table value with an alpha of 5% and df (n-1). The t-table value is (2.145). This indicates a significant difference between pre and post interventions of Acupressure and Foot Hydrotherapy using red ginger. The intervention resulted in a significant decrease in blood pressure in pregnant women with preeclampsia.

Conclusion

1. There is an effect of Acupressure and Foot Hydrotherapy using red ginger (*Zingiber officinale var rubrum*) on lowering blood pressure in pregnant women with preeclampsia at Puskesmas Kembaran I
2. There is a very strong correlation between Acupressure and Foot Hydrotherapy therapy using red ginger (*Zingiber officinale var rubrum*) in lowering blood pressure in pregnant women with preeclampsia at Puskesmas Kembaran I
3. The intervention or therapy of Acupressure and Foot Hydrotherapy using red ginger (*Zingiber*

officinale var rubrum) is effective in lowering blood pressure in pregnant women with preeclampsia

A. Suggestions

1. For the reduction of pre and post systolic, pre and post diastolic, and pre and post MAP blood pressure, pregnant women with preeclampsia should minimally undergo care according to health standards.
2. To lower blood pressure in pregnant women with preeclampsia, it is recommended to undergo Acupressure and Foot Hydrotherapy using red ginger (*Zingiber officinale var rubrum*).

B. For Future Researchers

1. Future researchers could expand this study by adding other research variables.
2. Future researchers should enrich the literature review on the combination of acupressure and hydrotherapy for pregnant women with preeclampsia, as there is still limited research on this variable

Acknowledgments

The author would like to express sincere gratitude to all institutions, organizations, and parties who have provided support for this research. Special thanks to the staff and management of Puskesmas Kembaran I for their cooperation and assistance in providing the necessary data for this study.

The author also deeply thanks the academic advisor at Poltekkes Kemenkes Semarang, who provided guidance, direction, and invaluable support throughout the research process. Last but not least, sincere thanks are also extended to all the respondents who agreed to participate in this study. Without their cooperation, this research would not have been possible. Thank you for all the support and encouragement provided.

References

1. Kementerian Kesehatan RI. Profil Kesehatan Indonesia 2023. Jakarta. 2024;
2. Dinas Kesehatan Kabupaten Banyumas. Laporan Kinerja Instansi Pemerintah Dinkes Banyumas Tahun 2023. Purwokerto; 2024 Mar.
3. Indrieni S. Asuhan Keperawatan Klien dengan Preeklampsi yang Di Rawat Di Rumah Sakit [Internet]. [Samarinda]: Politeknik Kesehatan Kementrian Kesehatan; 2020 [cited 2024 Apr

- 13]. Available from: <https://repository.poltekkes-kaltim.ac.id/1081/1/KTI%20SUSI%20INDRI%20ENI.pdf>
4. Obstetri P, Indonesia G, Kedokteran H, Maternal F. Pedoman Nasional Pelayanan Kedokteran Diagnosis dan Tata Laksana Pre-Eklamsia. 2016.
5. Kementerian Kesehatan RI. Buku KIA Kesehatan Ibu dan Anak [Internet]. Jakarta: Kementerian Kesehatan RI; 2023 [cited 2024 Apr 13]. Available from: <https://ayosehat.kemkes.go.id/pub/files/a8c1c20728a2d8d55f16a7e24f52cf97.pdf>
6. Puskesmas Kembaran I. Profil Kesehatan Puskesmas Kembaran I Tahun 2023. 2023;
7. Nisa atun, Choirin Nisa I, Faridasari I, Tinggi Ilmu Kesehatan Cirebon S. Pengaruh Akupresur terhadap Hipertensi pada Ibu Hamil Trimester II.
8. Widiastini L. Literature Review: Foot Massage dan Terapi Rendam Air Hangat Terhadap Edema Ekstremitas Bawah Pada Ibu Hamil. Jurnal Ilmiah Kebidanan (The Journal Of Midwifery). 2022 Nov 24;10(2):204–11.
9. Erlina Y, Dian Lestari Y, Studi III Keperawatan PD, Keperawatan Efarina Purwakarta AR, Barat J. Penerapan Terapi Rendam kaki dengan air Hangat (Hidroterapi) untuk menurunkan Tekanan Darah ypada IBu hamil Trimester III dengan Preeklamsia Ringa di Rumah Sakit Umum ASRI Purwakarta. 2023 [cited 2024 Apr 13];7(3). Available from: <https://journal.ummat.ac.id/index.php/jpmb/article/view/17253/7799>
10. Silfiyani L, Khayati N. Aplikasi foot hydrotheraphy dengan jahe merah (zingiber officinale var rubrum) terhadap penurunan tekanan darah pada lanjut usia dengan hipertensi. Ners Muda. 2021 Dec 31;2.
11. Setyaningsih S, Laxiana I. Aktivitas Antioksidan pada Permen Jelly dengan Bahan Baku Ekstrak rimpang Jahe. Bulletin Agroindustri [Internet]. 2022 Aug 29 [cited 2024 Apr 15];48(1):22–32. Available from: <https://intan.e-journal.id/agroindustri/article/view/19>
12. Susilowati. 50 Herbal Dan Suplemen Yang Memperpanjang Usia. Yogyakarta: Kyta; 2016.
13. Dahlan MS (2021a). Besar Sampel dan Cara Pengambilan Sampel dalam Penelitian Kedokteran dan Kesehatan [Internet]. 5th edn. Jakarta: Epidemiologi Indonesia, editor. Jakarta: Epidemiologi Indonesia; Available from: <http://www.penerbitsalemba.com>
14. Khairunnisa K, Sari FF, Anggelena M, Agustina D, Nursa'adah E. Penggunaan Effect Size Sebagai Mediasi dalam Koreksi Efek Suatu Penelitian. Jurnal Pendidikan Matematika (Judika Education). 2022 Dec 31;5(2):138–51.